ELSEVIER

Toxicology 120 (1997) 249–250

TOXICOLOGY

Volume contents

Volume 120 (1997)

In vitro interference with hepatocellular uptake of bile acids by xylene
M. Neghab, N.H. Stacey (Australia) 1–10
Renal transepithelial secretion of ochratoxin A in the non-filtering toad kidney
E. Bahnemann, H.-P. Kerling, S. Ensminger, G. Schwerdt, S. Silbernagl, M. Gekle (Germany) 11–17
Thymic regression and apoptosis in the rat after treatment with the Leydig cell cytotoxin ethylene dimethanesulphonate (EDS)
I.D. Morris, R.G. Lendon, C. Waters, G. Naylor, N. Jones (UK) 19–27
Cytotoxicity of food preservatives in cultured rat hepatocytes loaded with linolenic acid
N. Sugihara, K. Shimomichi, K. Furuno (Japan) 29–36
Loss of the metal binding properties of metallothionein induced by hydrogen peroxide and free radicals
I. Jiménez, M. Gotteland, A. Zarzuelo, R. Uauy, H. Speisky (Chile, Spain) 37–46
Outputs of hepatic copper and cadmium stimulated by tetrathiomolybdate (TTM) injection in Long-Evans Cinnamon (LEC) rats pretreated with cadmium, and in Fischer rats pretreated with copper and cadmium
Y.-R. Lai, N. Sugawara (Japan) 47–54
Nephrotoxic effects of bacterial ribonucleases in the isolated perfused rat kidney
O.N. Ilinskaya, S. Vamvakas (Russia, Germany) 55–63

Immunotoxicology
Assessment of the sensitization potential of five metal salts in the murine local lymph node assay
C. Mandervelt, F.L. Clottens, M. Demedts, B. Nemery (Belgium) 65–73

Corrigendum 75
Announcement 77

Antimitotic intervention with colchicine alters the outcome of *o*-DCB-induced hepatotoxicity in Fischer 344 rats
In Fischer 344 rats, higher rates of compensatory tissue repair may overcome more severe liver injury
S.G. Kulkarni, A. Warbritton, T.J. Bucci, H.M. Mehendale (USA) 79–88
Aluminum citrate is transported from brain into blood via the monocarboxylic acid transporter located at the blood-brain barrier
A study using microdialysis probes implanted into the jugular vein and cerebral cortex
D.C. Ackley, R.A. Yokel (USA) 89–97
Effects of low concentrations of 3-carene on alveolar macrophages in vitro
3-carene inhibits phagocytosis and dose-dependently kills the cells
A. Johansson, M. Lundborg (Sweden) 99–104
A model for quantitative measurement of sulfur mustard skin lesions in the rabbit ear
Epidermal thickness is a dose dependent measure of skin vesiculation
A. Zlotogorski, M. Goldenhersh, A. Shafran (Israel) 105–110
Changes in high molecular weight DNA fragmentation following human blood exposure to styrene-7,8-oxide
Styrene-7,8-oxide leads to decrease of total DNA fragments in white blood cells but not by direct action
B. Marczynski, M. Peel, X. Baur (Germany, Canada) 111–117

Kupffer cells from halothane-exposed guinea pigs carry trifluoroacetylated protein adducts
Kupffer cells may present altered proteins to the immune system
S.M. Furst, D. Luedke, A.J. Gandolfi (USA) 119–132
Comparative study of the damage produced by acute ethanol and acetaldehyde treatment in a human fetal hepatic cell line
The effects of direct toxicity of ethanol and acetaldehyde are demonstrated
I.P. Olivares, L. Bucio, V. Souza, A. Cárabez, M.C. Gutiérrez-Ruiz (Mexico) 133–144
Evaluation of the toxicity of ISIS 2302, a phosphorothioate oligonucleotide, in a four-week study in cynomolgus monkeys
Changes in clotting time and in the kidney seen at the higher doses were dose-dependent and reversible
S.P. Henry, H. Bolte, C. Auletta, D.J. Kornbrust (USA) 145–155
Announcement 157

Changes in neuroreceptor function of tracheal smooth muscle following acute ozone exposure of guinea pigs
Ozone may alter the responsiveness of tracheal lumen regulation
H.J.M. van Hoof, H.-P. Voss, K. Kramer, A.J.F. Boere, J.A.M.A. Dormans, L. van Bree, A. Bast (The Netherlands) 159–169
Polychlorinated biphenyl congener 153-induced ultrastructural alterations in rat liver: a quantitative study
Hepatocyte hypertrophy, increased SER profiles, abnormal mitochondria and augmentation of lysosomal elements and peroxisomes were found
J. Peng, A. Singh, W.P. Ireland, I. Chu (Canada) 171–183
Airborne particulate matter modulates the production of reactive oxygen species in human polymorphonuclear granulocytes
Aqueous and organic extracts differently affect chemoluminescence
B. Hitzfeld, K.H. Friedrichs, J. Ring, H. Behrendt (Germany) 185–195
Modulation of cytochrome P450 by 5,5′-bis-trifluoromethyl-2,2′-dichlorobiphenyl, a unique environmental contaminant
The compound has significant dioxin-like activity
A.T. Drahushuk, C.O. Choy, S. Kumar, J.H. McReynolds, J.R. Olson (USA). 197–205
A small peptide from durum wheat gliadin prevents cell agglutination induced by prolamin-peptides toxic in coeliac disease
The in vitro technique shows interesting antogonisms between related proteins
M. De Vincenzi, G. Gasbarrini, V. Silano (Italy) 207–213
Cadmium uptake by a human hepatic cell line (WRL-68 cells)
Uptake pathways appear in human hepatocytes appear to be similar to those in rat hepatocytes
V. Souza, L. Bucio, M.C. Gutiérrez-Ruiz (Mexico). 215–220
Effect of fructose on the biochemical toxicity of hydrazine in isolated rat hepatocytes
10 mM fructose ameliorates three parameters for cytotoxicity
M.D.J. Kerai, J.A. Timbrell (UK) 221–230

Immunotoxicology
Prenatal exposure to cypermethrin modulates rat NK cell cytotoxic functions
Prenatal exposure of rats to cypermethrin resulted in decreased NK cell and antibody-dependent cytotoxicity activity
G. Santoni, F. Cantalamessa, L. Mazzucca, S. Romagnoli, M. Piccoli (Italy) 231–242